

For Research Use Only

RNF40 Recombinant antibody

Catalog Number: 84058-4-RR



Basic Information

Catalog Number: 84058-4-RR	GenBank Accession Number: BC006133	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 9810	CloneNo.: 241214D12
Source: Rabbit	UNIPROT ID: O75150	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000
Isotype: IgG	Full Name: ring finger protein 40	
Immunogen Catalog Number: AG8047	Calculated MW: 114 kDa	
	Observed MW: 130-140 kDa	

Applications

Tested Applications:
WB, IHC, ELISA

Species Specificity:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, mouse brain tissue, A375 cells, HEK-293 cells, HepG2 cells, MCF-7 cells, mouse liver tissue

IHC: mouse testis tissue,

Background Information

The DNA damage response (DDR) is emerging as a vast signaling network that temporarily modulates numerous aspects of cellular metabolism in the face of DNA lesions, especially critical ones such as the double-strand break (DSB). The DDR involves extensive dynamics of protein post-translational modifications, most notably phosphorylation and ubiquitylation. A recently identified novel component of the DDR pathway - histone H2B monoubiquitylation - exemplifies this principle. Monoubiquitination of histone H2B is a dynamic post-translational histone modification associated with transcriptional elongation and the DNA damage response [PMID:22021426]. In mammalian cells, H2B monoubiquitylation is driven primarily by an E3 ubiquitin ligase composed of the two RING finger proteins RNF20 and RNF40 [PMID:22031019,21827756].

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84058-4-RR (RNF40 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 84058-4-RR (RNF40 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 84058-4-RR (RNF40 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 84058-4-RR (RNF40 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).