

For Research Use Only

# IDO1 Recombinant antibody

Catalog Number: 84063-1-RR



## Basic Information

<b>Catalog Number:</b> 84063-1-RR	<b>GenBank Accession Number:</b> BC027882	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1000 µg/ml	<b>GeneID (NCBI):</b> 3620	<b>CloneNo.:</b> 241110C8
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P14902	<b>Recommended Dilutions:</b> WB 1:1000-1:8000
<b>Isotype:</b> IgG	<b>Full Name:</b> indoleamine 2,3-dioxygenase 1	
<b>Immunogen Catalog Number:</b> AG3953	<b>Calculated MW:</b> 403 aa, 45 kDa	
	<b>Observed MW:</b> 45 kDa	

## Applications

<b>Tested Applications:</b> WB, FC (Intra), ELISA	<b>Positive Controls:</b> WB : IFN gamma treated HeLa cells,
<b>Species Specificity:</b> human	

## Background Information

IDO1 is the target for therapy in a range of clinical settings, including cancer, chronic infections, autoimmune and allergic syndromes, and transplantation. Elevated IDO1 expression is a hallmark of major viral infections including HIV, HBV, HCV or influenza and also of major bacteria infections, such as Tb, CAP, listeriosis and sepsis. Pathogens are able to hijack the immunosuppressive effects of IDO1 and make use of them to facilitate their own life cycle. MW of IDO1 is 40-42kd (PMID: 14502282; 17055065).

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data



IFN gamma treated HeLa cells were subjected to SDS PAGE followed by western blot with 84063-1-RR (IDO1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



$1 \times 10^6$  IFN-gamma treated HeLa cells were intracellularly stained with 0.25  $\mu$ g Ido1 Recombinant Antibody (84063-1-RR, Clone:241110C8) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25  $\mu$ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Bi-layer interferometry (BLI) kinetic assays of 84063-1-RR against Human IDO1 were performed. The affinity constant is below 1 pM.