## For Research Use Only

## Leptin Recombinant antibody

Catalog Number:84072-5-RR



**Basic Information** 

Catalog Number:

84072-5-RR

Size: 1000 μg/ml

Source: Rabbit BC060830 GeneID (NCBI):

GenBank Accession Number:

3952 UNIPROT ID: P41159

Isotype: Full Name: IgG leptin

Calculated MW: 167 aa, 19 kDa

**Applications** 

Tested Applications: IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Purification Method: Protein A purfication

CloneNo.:

IF/ICC 1:125-1:500

240720A6

Recommended Dilutions:

Positive Controls: IF/ICC: 3T3-L1 cells,

## **Background Information**

Leptin is a protein that is secreted by white adipocytes, and which plays a major role in the regulation of body weight. Leptin is the most critical hormone in the homeostatic regulation of energy balance among those so far discovered. Leptin primarily acts on the neurons of the mediobasal part of hypothalamus to regulate food intake, thermogenesis, and the blood glucose level. Leptin also has several endocrine functions, and is involved in the regulation of immune and inflammatory responses, hematopoiesis, angiogenesis and wound healing. Mutations in this gene and/or its regulatory regions cause severe obesity, and morbid obesity with hypogonadism. Leptin has also been linked to type 2 diabetes mellitus development.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunofluorescent analysis of (4% PFA) fixed 3T3-L1 cells using LEP antibody (84072-5-RR, Clone: 240720A6) at dilution of 1:250 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug LEP Recombinant antibody (84072-5-RR, Clone:240720A6) and APC-Conjugated Goat Anti-Rabbit I gG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug LEP Recombinant antibody (84072-5-RR, Clone:240720A6) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 84072-5-RR against Human LEP were performed. The affinity constant is 35.3 nM.