

For Research Use Only

MYO6 Recombinant antibody

Catalog Number: 84085-3-RR



Basic Information

Catalog Number:

84085-3-RR

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG24906

GenBank Accession Number:

BC146764

GeneID (NCBI):

4646

UNIPROT ID:

Q9UM54

Full Name:

myosin VI

Calculated MW:

1285 aa, 149 kDa

Observed MW:

145-150 kDa

Purification Method:

Protein A purification

CloneNo.:

241251G11

Recommended Dilutions:

WB 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : MCF-7 cells, LNCaP cells, HEK-293T cells, DU 145 cells

Background Information

MYO6, an actin-based motor protein, is the only myosin known to move toward the minus end of actin filaments. MYO6 is highly expressed in the inner and outer hair cells of the ear, retina, and polarized epithelial cells such as kidney proximal tubule cells and intestinal enterocytes. And it participates in a wide range of biological processes within cells, including clathrin-mediated endocytosis, vesicular membrane traffic, polarized secretion, and autophagy (PMID: 23620821; PMID: 28591580). Previous studies showed that MYO6 is upregulated in various types of cancer, and it has been widely reported to contribute to tumor cell migration and metastasis. Some articles indicate that MYO6 is associated with prostate cancer, lung cancer, human colorectal cancer and gastric cancer (PMID: 29022908).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84085-3-RR (MYO6 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



LNCaP cells were subjected to SDS PAGE followed by western blot with 84085-3-RR (MYO6 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.