

For Research Use Only

PDCD4 Recombinant antibody

Catalog Number: 84162-3-RR



Basic Information

Catalog Number: 84162-3-RR	GenBank Accession Number: BC026104	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 27250	CloneNo.: 241463E3
Source: Rabbit	UNIPROT ID: Q53EL6	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: programmed cell death 4 (neoplastic transformation inhibitor)	
Immunogen Catalog Number: AG3288	Calculated MW: 469 aa, 52 kDa	
	Observed MW: 54-64 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : BxPC-3 cells, HeLa cells, HUVEC cells, MCF-7 cells, HEK-293 cells
Species Specificity: human	

Background Information

Programmed cell death 4 (Pdc4) is a novel tumor suppressor that inhibits translation, progression and invasion. It was first identified as being differentially upregulated during apoptosis. Pdc4 interferes with the activity of the eukaryotic initiation factor (eIF) 4A by displacing the scaffold protein eIF4G from its binding to the RNA helicase eIF4A. The calculated molecular weight of unmodified PDCD4 protein is 52 kDa, but the modified protein is about 54-64 kDa.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84162-3-RR (PDCD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 HeLa cells were intracellularly stained with 0.25 μ g PDCD4 Recombinant antibody (84162-3-RR, Clone:241463E3) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Bioluminescence resonance energy transfer (BRET) kinetic assays of 84162-3-RR against Human PDCD4 were performed. The affinity constant is 1.54 nM.