For Research Use Only

## PCSK9 Recombinant antibody

Catalog Number:84172-4-RR





Basic Information	Catalog Number: 84172-4-RR	GenBank Accession Number: NM_174936.4	Purification Method: Protein A purification
	Size: 1000 µg/ml	GenelD (NCBI): 255738	CloneNo.: 241395G1
	Source: Rabbit	UNIPROT ID: Q8NBP7	Recommended Dilutions: WB 1:500-1:2000
	Isotype: IgG	Full Name: proprotein convertase subtilisin/kexin type 9	
		Calculated MW: 74 kDa	
		Observed MW: 72-78 kDa, 62 kDa	
Applications	Tested Applications:	Positive	Controls:
	WB, ELISA WB : HepG2 cells, Species Specificity: human		G2 cells,
Background Information	Proprotein convertase subtilisin/kexin type 9 (PCSK9) is a crucial protein governing the circulating levels of low density lipoprotein-cholesterol (LDL-C), by virtue of its pivotal role in the degradation of the LDL receptor (LDLR). PCSK9 is expressed in the kidney and lung. It is synthesized as a 72 kDa immature precursor that undergoes autocatalytic cleavage in the endoplasmic reticulum to generate a 63 kDa mature protein. The cleaved N-terminal fragment remains associated with the mature protein and is necessary for its secretion, allowing it to circulate in the blood. The ability of PCSK9 to regulate a diverse group of cell-surface proteins hinted that it might also be able to influence additional membrane proteins that are important in anti-tumour immune responses. Targeting PCSK9 to treat cancer is also attractive because two neutralizing antibodies against it, evolocumab and alirocumab, have already been approved for human clinical use to lower cholesterol levels. (PMID: 30522786, PMID: 22493497)		
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azide Aliquoting is unnecessary for	e and 50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





HepG2 cells were subjected to SDS PAGE followed by western blot with 84172-4-RR (PCSK9 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Biolayer interferometry (BLL) kinetic assays of 84172-4-RR against Human PCSK9 were performed. The affinity constant is 0.80 nM.