For Research Use Only

Ki-67 Recombinant antibody, PBS Only

Catalog Number:84192-4-PBS




: CloneNo.: 241499E7 fied by monoclonal	
fied by monoclonal	
<i>I</i> :	
marker for proliferation. Ki67 is present during all active nt in resting cells (GO). Cellular content of Ki-67 protein 5 phase of the cell cycle. Therefore, the nuclear expression of ny immunohistochemistry. It has been demonstrated to be of ancer, several studies have reported an association between	
tore it immediately at -80°C	
S b	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84192-4-RR (Ki-67 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84192-4-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84192-4-RR (Ki-67 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84192-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human lung cancer tissue using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84192-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84192-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human lung cancer tissue using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:200 and Multi-rAb CoraLite © Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84192-4-PBS in a different storage buffer formulation.