For Research Use Only

Ki-67 Recombinant antibody

Catalog Number:84192-4-RR



Basic Information

Catalog Number: GenBank Accession Number: 84192-4-RR NM_002417

Size: GeneID (NCBI): 1000 μ g/ml 4288

Rabbit P46013

Isotype: Full Name: antigen identified by monoclonal

antibody Ki-67 Calculated MW: 359 kDa

UNIPROT ID:

Purification Method:

Protein A purfication CloneNo.:

241499E7 Recommended Dilutions: IHC 1:250-1:1000

IF-P 1:50-1:500 IF/ICC 1:200-1:800

Applications

Tested Applications: IHC, IF/ICC, IF-P, ELISA Species Specificity:

human

Source:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human tonsillitis tissue, human Breast cancer tissue, human malignant melanoma tissue, human prostate cancer tissue, Human Rectal cancer tissue, Human renal cancer, human skin cancer tissue, Human Spleen tissue

IF-P: human lung cancer tissue, human thyroid cancer tissue

IF/ICC : HeLa cells, MCF-7 cells, U2OS cells, HepG2 cells, A431 cells, A549 cells

Background Information

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84192-4-RR (Ki-67 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84192-4-RR (Ki-67 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human lung cancer tissue using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:200 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human lung cancer tissue using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).