For Research Use Only

CD3 Delta Recombinant antibody

Catalog Number:84214-4-RR



Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:500-1:2000

CloneNo.:

241523H2

Basic Information

Catalog Number: GenBank Accession Number: 84214-4-RR BC070321

Size: Genel D (NCBI): 1000 ug/ml 915

Source: UNIPROT ID: Rabbit P04234
Isotype: Full Name:

IgG CD3d molecule, delta (CD3-TCR

Calculated MW: 171 aa, 19 kDa Observed MW: 20-25 kDa

complex)

Applications

Tested Applications: WB, IHC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Jurkat cells, MOLT-4 cells IHC: human tonsillitis tissue,

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is complex, leading to the intracellular signal transduction. CD3 is complex per lemmbrane protein which consists of an extracellular domain of 84 amino acids, a transmembrane region, and a cytoplasmic domain of 45 amino acids. Defects in CD3 delta are a cause of severe combined immunodeficiency.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84214-4-RR (CD3 Delta antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84214-4-RR (CD3 Delta antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84214-4-RR (CD3 Delta antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).