

For Research Use Only

VDAC2 Recombinant antibody, PBS Only

Catalog Number: 84225-3-PBS



Basic Information

Catalog Number: 84225-3-PBS	GenBank Accession Number: BC000165	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 7417	CloneNo.: 241569C3
Source: Rabbit	UNIPROT ID: P45880	
Isotype: IgG	Full Name: voltage-dependent anion channel 2	
Immunogen Catalog Number: AG2266	Calculated MW: 294 aa, 32 kDa	

Applications

Tested Applications:
IHC, IF/ICC, FC (Intra), Indirect ELISA

Species Specificity:
human, mouse

Background Information

VDACs (Voltage Dependent Anion selective Channels), also known as mitochondrial porins, are a family of pore-forming proteins discovered in the mitochondrial outer membrane. Mammals show a conserved genetic organization of the VDAC genes. It's reported that the amount of VDAC transcripts in liver is usually lower than in the other tissues. VDAC2 and especially VDAC3 are highly expressed in testis, while mouse VDAC1 is poorly expressed in this tissue. (PMID: 22020053)

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using VDAC2 antibody (84225-3-RR, Clone: 241569C3) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 84225-3-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 84225-3-RR (VDAC2 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84225-3-PBS in a different storage buffer formulation.



1x10⁶ HeLa cells were intracellularly stained with 0.25 ug Vdac2 Recombinant Antibody (84225-3-RR, Clone:241569C3) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84225-3-PBS in a different storage buffer formulation.