For Research Use Only

Siglec-7/CD328 Recombinant antibody, proteintech® PBS Only


www.ptglab.com


Catalog Number:84253-4-PBS


Basic Information	Catalog Number: 84253-4-PBS	GenBank Accession Number: NM_014385.4	Purification Method: Protein A purfication
	Size: 1 mg/ml	Genel D (NCBI): 27036	CloneNo.: 241525G3
	Source: Rabbit	UNIPROT ID: Q9Y286-1	
	Isotype: IgG	Full Name: sialic acid binding Ig-like lectin 1	7
		Calculated MW: 51 kDa	
		Observed MW: 75 kDa	
Applications	Tested Applications: WB, IHC, Indirect ELISA		
	Species Specificity: human		
Background Information	Sialic acid binding Ig-like lectin 7 (Siglec-7), also known as CD328 or p75/AIRM-1, is a member of the Siglec family of glycan-recognition proteins. Siglec-7 is a type-I transmembrane protein consisting of three extracellular immunoglobulin-like domains that comprise an N-terminal V-set domain and two C2-set domains, a transmembrane region and a cytoplasmic tail containing two tyrosine residues embodied in immunoreceptor tyrosine-based inhibition motif-like motifs (PMID: 32322597; 10567377). It is mainly expressed on immune cells, with low levels on granulocytes, intermediate levels on monocytes, and relatively high levels on a major subset of CD8+T cells (PMID: 10567377). Siglec-7 is an inhibitory receptor that negatively regulates the function of NK cells and modulates the immune response through the interaction of sialic acid-containing ligands (PMID: 27312286).		
Storage	Storage: Store at -80°C. The product is shipped with Storage Buffer: PBS Only	ice packs. Upon receipt, store it immediatel	yat-80°C

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data








U-937 cells were subjected to SDS PAGE followed by western blot with 84253-4-RR (Siglec-7/CD328 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84253-4-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84253-4-RR (Siglec-7/CD328 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84253-4-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84253-4-RR (Siglec-7/CD328 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84253-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 84253-4-RR against Human Siglec-7/CD328 were performed. The affinity constant is 53.4 nM.