For Research Use Only

GBP2 Recombinant antibody

Catalog Number:84264-6-RR




Basic Information	Catalog Number: 84264-6-RR	GenBank Accession Number: BC022272	Purification Method: Protein A purfication	
	Size: 1000 µg/ml	GenelD (NCBI): 2634	CloneNo.: 241580C12	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number:	UNIPROT ID: P32456 Full Name: guanylate binding protein 2, interferon-inducible	Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:125-1:500	
				AG2425
		Observed MW: 67 kDa		
	Applications	Tested Applications:	F/ICC, FC (Intra), ELISA WB : A549 cells, IM-9 cells, HaCaT cells, K-562 cells ies Specificity: IF/ICC : HeLa cells.	
WB, IF/ICC, FC (Intra), ELISA Species Specificity: human				
Background Information	marker of interferon (IFN) respons stimulation. Increased GBP2 expr role in T-cell defense against brea activation and matrix metallopro	GBP2 is a member of the GTPase family and is crucial to host immunity against pathogens. GBP2 is also used as a marker of interferon (IFN) responsiveness because it is one of the most highly expressed genes after IFN- γ stimulation. Increased GBP2 expression is also associated with a better prognosis in breast cancer and may have a role in T-cell defense against breast cancer. Interestingly, GBP2 expression in NIH/3T3 fibroblasts inhibited Rac activation and matrix metalloproteinase-9 expression, suggesting a possible role for GBP2 in regulating cancer metastasis. The molecular weight of GBP2 is 67 kDa. (PMID: 29072687, 35383115)		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WWW.PTGLAB.COM

Various lysates were subjected to SDS PAGE followed by western blot with 84264-6-RR (GBP2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using GBP2 antibody (84264-6-RR, Clone: 241580C12) at dilution of 1:250 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 A549 cells were intracellularly stained with 0.2 ug GBP2 Recombinant antibody (84264-6-RR, Clone:241580C12) and Multi-rAb CoraLite ® Plus 647-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR005)(red), or 0.2 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 84264-6-RR against Human GBP2 were performed. The affinity constant is below 1 pM.