For Research Use Only

CD72 Recombinant antibody

Catalog Number:84270-5-RR



Purification Method:

Protein A purfication

WB 1:1000-1:4000

Recommended Dilutions:

CloneNo.:

241301B3

Basic Information

Catalog Number:

84270-5-RR

Size: 1000 μg/ml

Source: Rabbit Isotype:

Calculate

359 aa, 40 kDa

Full Name:

BC030227

GeneID (NCBI):

UNIPROT ID: P21854

GenBank Accession Number:

CD72 molecule
Calculated MW:

Applications

Tested Applications:

WB, ELISA

Species Specificity: human

Positive Controls:

WB: Raji cells,

Background Information

CD72 is a type II transmembrane glycoprotein of 39-45 kDa and is expressed as a disulphide-linked homodimer (PMID: 1711157). CD72 is a pan B-cell marker covering the full range of differentiation from the very early stages of B cells to mature surface Ig+ B cells (PMID: 1384316). It is expressed in all stages of B cell development except plasma cells. CD72 is also expressed by dendritic cells, tissue macrophages in the red pulp of the spleen and von Kupffer cells in the liver (PMID: 8561576; 1384316). CD72 is the ligand for CD5 (PMID: 1711157). CD72 is a negative regulator of BCR signaling (PMID: 9590210). Interaction of CD100 with CD72 strictly tunes the strength of BCR signals and is essential for maintaining immunological homeostasis as well as generating a proper immune response (PMID: 16113236).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Raji cells were subjected to SDS PAGE followed by western blot with 84270-5-RR (CD72 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 84270-5-RR against Human CD72 were performed. The affinity constant is 59.3 pM.