

For Research Use Only

# CD44 Recombinant antibody

Catalog Number: 84369-5-RR



## Basic Information

|                                      |  |   |
|--------------------------------------|--|---|
| <b>Catalog Number:</b><br>84369-5-RR | <b>GenBank Accession Number:</b><br>XM_006234631.3 | <b>Purification Method:</b><br>Protein A purification                 |
| <b>Size:</b><br>1000 µg/ml           | <b>GeneID (NCBI):</b><br>25406                     | <b>CloneNo.:</b><br>241648D1  |
| <b>Source:</b><br>Rabbit             | <b>UNIPROT ID:</b><br>O70509                       | <b>Recommended Dilutions:</b><br>WB 1:1000-1:4000<br>IHC 1:400-1:1600 |
| <b>Isotype:</b><br>IgG               | <b>Full Name:</b><br>Cd44 molecule                 |   |
|                                      | <b>Calculated MW:</b><br>40 kDa                    |   |
|                                      | <b>Observed MW:</b><br>80 kDa                      |   |

## Applications

|   |   |
|---|---|
| <b>Tested Applications:</b><br>WB, IHC, ELISA | <b>Positive Controls:</b>   |
| <b>Species Specificity:</b><br>rat            | <b>WB :</b> rat spleen tissue,<br><b>IHC :</b> rat spleen tissue, |

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

CD44 is a type I transmembrane glycoprotein expressed on embryonic stem cells and in various levels on other cell types including connective tissues and bone marrow. CD44 expression is also upregulated in subpopulations of cancer cells and is recognized as a molecular marker for cancer stem cells (PMID: 29747682). It is a cell-surface receptor that mediates cell-cell and cell-matrix interactions through its affinity for hyaluronic acid (HA) and possibly also through its affinity for other ligands (PMID: 10694938). Adhesion with HA plays an important role in cell migration, tumor growth and progression. CD44 is also involved in lymphocyte activation, recirculation and homing, and in hematopoiesis.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




rat spleen tissue was subjected to SDS PAGE followed by western blot with 84369-5-RR (CD44 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded rat spleen tissue slide using 84369-5-RR (CD44 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat spleen tissue slide using 84369-5-RR (CD44 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 84369-5-RR against Rat CD44 were performed. The affinity constant is below 1 pM.