

For Research Use Only

# Transferrin/TF Recombinant antibody, PBS Only

Catalog Number:84386-5-PBS



## Basic Information

<b>Catalog Number:</b> 84386-5-PBS	<b>GenBank Accession Number:</b> BC059367	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1 mg/ml	<b>GeneID (NCBI):</b> 7018	<b>CloneNo.:</b> 241769C8
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P02787	
<b>Isotype:</b> IgG	<b>Full Name:</b> transferrin	
	<b>Calculated MW:</b> 698 aa, 77 kDa	
	<b>Observed MW:</b> 77 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
human, mouse, rat

## Background Information

Serotransferrin(TF) is a 77 kDa secreted protein which is also named as transferrin, siderophilin, beta-1 metal-binding globulin and belongs to a family of homologous iron-binding glycoproteins that encompasses lactoferrin (found both intracellular and in secretions, including milk), melanotransferrin (present on melanoma cells) and ovotransferrin (present in egg white). TF is a multi-function protein with a primary role in transporting iron in a safe, redox-inactive state from absorption to utilization or storage sites around the body. The association of Tf with the immune system derives from its ability to restrict serum free-iron levels, creating low-iron environments where the infection capacity of pathogenic microorganisms is limited. TF is expressed predominantly in the liver and secreted in plasma, but lower amounts can be synthesized in other tissues such as the brain and the testis.

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84386-5-RR (TF antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84386-5-PBS in a different storage buffer formulation.



human urine sample were subjected to SDS PAGE followed by western blot with 84386-5-RR (TF antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84386-5-PBS in a different storage buffer formulation.