

For Research Use Only

IL-10RA Recombinant antibody

Catalog Number: 84428-3-RR



Basic Information

Catalog Number:

84428-3-RR

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC028082

GeneID (NCBI):

3587

UNIPROT ID:

Q13651

Full Name:

interleukin 10 receptor, alpha

Calculated MW:

578 aa, 63 kDa

Observed MW:

90 kDa

Purification Method:

Protein A purification

CloneNo.:

241833D5

Recommended Dilutions:

WB 1:200-1:1000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : HepG2 cells, K-562 cells

Background Information

Interleukin-10 (IL-10) is an anti-inflammatory cytokine that plays a critical role in maintaining immune system balance. IL-10 signals through a receptor complex consisting of two molecules of the IL-10R alpha-chain (IL-10RA) and two molecules of the accessory IL-10R beta-chain (IL-10RB) (PMID: 22236434). IL-10RA and IL-10RB are members of the type II cytokine receptor family. IL-10RA, also known as IL-10R1 or CD210, is expressed on most hematopoietic cells at a basal level but is upregulated by various cells upon activation (PMID: 24507158). IL-10RA serves as the ligand-binding subunit of the receptor complex. Upon binding of IL-10 to the IL-10R complex, the kinases JAK1 and TYK2 are activated and catalyze the phosphorylation of IL-10RA at specific intracellular tyrosine residues, leading to the recruitment and subsequent phosphorylation of STAT3 (PMID: 11244051; 10433356).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84428-3-RR (IL-10RA antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.