For Research Use Only

RBP4 Recombinant antibody

Catalog Number:84496-4-RR




Basic Information	Catalog Number: 84496-4-RR	GenBank Accession Number: BC020633	Purification Method: Protein A purification
	Size: 1000 µg/ml	GeneID (NCBI): 5950	CloneNo.: 241944H4
	Source: Rabbit	UNIPROT ID: P02753	Recommended Dilutions: WB 1:5000-1:50000
	lsotype: IgG	Full Name: retinol binding protein 4, plasma Calculated MW: 201 aa, 23 kDa	
		Applications	Tested Applications: WB, ELISA Species Specificity: human
WB : HepC	52 cells, human plasma ، human liver tissue		
Background Information	RBP4 (retinol-binding protein 4) is a carrier protein that transports vitamin A (retinol) from the liver to the periphera tissues. Synthesized primarily by hepatocytes and adipocytes as a 21 kDa non-glycosylated protein, RBP4 is secreted into the circulation as a retinol-RBP4 complex. In plasma the RBP4-retinol complex is bound to transthyretin (TRR), which prevents prevent kidney filtration. Two truncated forms of RBP4, RBP4-L (truncated at Leu-183) and RBP4-LL (truncated at Leu-182 and Leu-183), exist by proteolytic process. RBP4-L and RBP4-LL, which do not bind TTR, are normally excreted into the urine but accumulate in the serum during renal failure. Urinary RBP4 has been reported as marker for glomerular disease. RBP4 also was identified as an adipokine that elevated in some INS-resistant states. Measurement of serum RBP4 could be used to assess the risk of INS resistance, type 2 diabetes, obesity, and cardiovascular disease. (18752671, 16034410)		
	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84496-4-RR (RBP4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Biolayer interferometry (BLI) kinetic assays of 84496-4-RR against Human RBP4 were performed. The affinity constant is 32.4 nM.

200 nM 100 nM 50 nM 25 nM 12.5 nM 6.25 nM 3.125 nM

400