For Research Use Only

## GLAST/EAAT1 Recombinant antibody, PBS Only




Catalog Number:84497-4-PBS

Basic Information	Catalog Number: 84497-4-PBS	GenBank Accession Number: BC037310	Purification Method: Protein A purfication
	Size: 1 mg/ml	GeneID (NCBI): 6507	CloneNo.: 241907H3
	Source: Rabbit	UNIPROT ID: P43003	
	Isotype: IgG Immunogen Catalog Number: AG16962	Full Name: solute carrier family 1 (glial high	
		affinity glutamate transporter), member 3	
		Calculated MW: 542 aa, 60 kDa	
Applications	Tested Applications: IF/ICC, FC (Intra), Indirect ELISA		
	Species Specificity: human, mouse		
Background Information	SLC 1A3, also known as EAAT-1 or GLAST, is a membrane-bound protein localized in glial cells and pre-synaptic glutamatergic nerve endings. It transports the excitatory neurotransmitters L-glutamate and D-aspartate, which is essential for terminating the postsynaptic acction of glutamate. Recently, a correlation between expression/function of glial EAAT-1 and tumor proliferation has been reported. The exceptionally rare expression of EAAT-1 in non-neoplastic choroid plexus (CP) compared to choroid plexus tumors (CPT) may distinguishes neoplastic from normal CP. There are a number of splicing variants of SLC1A3, like GLAST1a and GLAST1b, exist due to the exon skipping. It also undergo glycosylation. Variety of bands can be observed in the western blotting assay: 50-55 kDa represents the unglycosylated GLAST1a or GLAST1b, 65-70 kDa correspond to the glycosylated proteins, larger proteins between 90-130 kDa may be the multimers of SLC1A3. (11086157, 17471058, 12546822)		
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS Only		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using GLAST antibody (84497-4-RR, Clone: 241907H3) at dilution of 1:250 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 84497-4-PBS in a different storage buffer formulation.



1x10^6 HEK-293 cells were intracellularly stained with 0.25 ug GLAST Recombinant antibody (84497-4-RR, Clone:241907H3) and APC-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84497-4-PBS in a different storage buffer



1x10^6 U-937 cells were intracellularly stained with 0.25 ug GLAST Recombinant antibody (84497-4-RR, Clone:241907H3) and APC-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84497-4-PBS in a different storage buffer