For Research Use Only

CD6 Recombinant antibody

Catalog Number:84508-5-RR



Purification Method:

IHC: human tonsillitis tissue,

Basic Information

Catalog Number: GenBank Accession Number: 84508-5-RR BC033755

 84508-5-RR
 BC033755
 Protein A purfication

 Size:
 GeneID (NCBI):
 CloneNo.:

 1000 μ g/ml
 923
 241965D11

 Source:
 UNIPROT ID:
 Recommended Dilutions:

 Rabbit
 P30203
 WB 1:500-1:2000

 Isotype:
 Full Name:
 IHC 1:500-1:2000

 IgG
 CD6 molecule

Observed MW: 140 kDa

Applications

Tested Applications: Positive Controls: WB, IHC, ELISA WB: Jurkat cells,

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

CD6 is a type I transmembrane glycoprotein that belongs to the scavenger receptor cysteine-rich superfamily (PMID: 21880988). It is composed of an extracellular region, which consists of three tandem SRCR domains, a transmembrane region, and a cytoplasmic tail devoid of intrinsic catalytic activity but harboring several phosphorylatable residues suitable for intracellular signal transduction (PMID: 38139340; 23711376). CD6 is expressed by T cells, medullary thymocytes, a subset of B cells and NK cells, and in some cells of the brain (PMID: 1919444; 21178331). CD6 plays a role in lymphocyte activation, proliferation, and survival processes via interaction with its endogenous ligands.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 84508-5-RR (CD6 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 percent. Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84508-5-RR (CD6 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84508-5-RR (CD6 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).