For Research Use Only

ALCAM Recombinant antibody

Catalog Number:84528-1-RR



Basic Information

Catalog Number:

GenBank Accession Number: NM_009655.3

Purification Method:

84528-1-RR

(LD (NCD))

Protein A purification

Size: 1000 μg/ml GeneID (NCBI): 11658 CloneNo.: 241945A4

Source:

UNIPROT ID:

Recommended Dilutions:

Rabbit Isotype:

Q61490 Full Name: WB 1:2000-1:10000 IHC 1:250-1:1000

IgG

activated leukocyte cell adhesion

molecule

Calculated MW:

65kDa

Observed MW:

90-100 kDa

Applications

Tested Applications:

Positive Controls:

WB, IHC, ELISA

WB: mouse lung tissue,

IHC: mouse liver tissue,

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate

buffer pH 6.0

Background Information

Activated leukocyte cell adhesion molecule (ALCAM, also known as CD166) is a cell adhesion molecule that belongs to the immunoglobulin superfamily. It is involved in cell-cell adhesion through homophilic and heterophilic (to CD6) interactions. ALCAM is widely expressed in a variety of normal tissues. Altered ALCAM expression has been associated with the differentiation state and progression in some neoplasms including melanoma, prostate, colorectal, and breast cancers (PMID: 20461761; 18172759).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse lung tissue was subjected to SDS PAGE followed by western blot with 84528-1-RR (ALCAM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 84528-1-RR (ALCAM antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 84528-1-RR (ALCAM antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).