## For Research Use Only

## FATP5/SLC27A5 Recombinant antibody

Catalog Number:84557-1-RR



**Purification Method:** 

Protein A purfication

**Recommended Dilutions:** 

WB 1:2000-1:10000

IHC 1:200-1:800

CloneNo.:

241884H2

**Basic Information** 

Catalog Number: GenBank Accession Number:

84557-1-RR BC148807 GeneID (NCBI): Size: 1000 µg/ml 10998 Source: **UNIPROT ID:** Q9Y2P5 Rabbit

Isotype: Full Name: solute carrier family 27 (fatty acid

transporter), member 5 Immunogen Catalog Number:

AG18276 Calculated MW:

> 690 aa, 75 kDa Observed MW: 70-75 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, ELISA Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed

with citrate buffer pH 6.0

Positive Controls:

WB: mouse liver tissue, rat liver tissue

IHC: mouse liver tissue.

## **Background Information**

SLC27A5, also known as FATP5, ACSVL6, and BAL, is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS) and BAL, is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS) and BAL, is an isozyme of very long-chain acyl-CoA synthetase (VLCS). It is an isozyme of very long-chain acyl-CoA synthetase (VLCS) and BAL, is an isozyme of very long-chain acyl-CoA synthetase (VLCS) and BAL, is an isozyme of very long-chain acyl-CoA synthetase (VLCS) and acyl-CoA synthetase (VLCS) andcapable of activating very long-chain fatty acids containing 24- and 26-carbons (PMID: 10479480). FATP5 mediates the import of long-chain fatty acids (LCFA) by facilitating their transport across cell membranes (PubMed:20448275). FATP5 is associated with endoplasmic reticulum but not with peroxisomes, and its primary role is in fatty acid elongation or complex lipid synthesis. FATP5 is exclusively expressed by the liver and localized to the basal plasma membrane of hepatocytes (PMID:16618416).


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 84557-1-RR (SLC27A5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 84557-1-RR (FATP5/SLC27A5 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 84557-1-RR (FATP5/SLC27A5 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).