For Research Use Only

CEBP Alpha/CEBPA Recombinant antibody, PBS Only (Detector)

Catalog Number:84668-2-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

84668-2-PBS

GeneID (NCBI):

Size: 1 mg/ml NM_004364 1050

CloneNo.: 242101C10

Source: Rabbit

UNIPROT ID: P49715 Full Name:

Isotype:

CCAAT/enhancer binding protein

Immunogen Catalog Number:

(C/EBP), alpha

AG29947

Calculated MW: 38 kDa

Observed MW: 40-45 kDa

Applications

Tested Applications:

WB, FC (Intra), Cytometric bead array, Indirect ELISA

Species Specificity:

Background Information

CEBPA and its isoforms play important roles in lineage determination and gene activation in a variety of cell types $by\ activating\ transcription\ from\ lineage-specific\ promoters.\ CEBPA\ is\ a\ DNA-binding\ protein\ that\ recognizes\ two$ different motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers. In hematopoiesis, C/EBPa is a key factor in driving the development of myeloid cells interacting with a variety of factors, including c-Myc, PU.1, and microRNAs. It can also form heterodimers with the related proteins CEBP-beta and CEBP-gamma. The encoded protein has been shown to bind to the promoter and modulate the expression of the gene encoding leptin which plays an important role in body weight homeostasis. CEBPA can interact with CDK2 and CDK4, thereby inhibiting these kinases and causing growth arrest in cultured cells. Several pathways have been implicated as the means by which CEBPA mediates cell cycle arrest and proliferation, including p21, cyclin-dependent kinases and the E2F complex via c-Myc. The calcualted molecular weight of CEBPA is 38 kDa, but modified CEBPA is about 42 kDa (PMID: 19623175).

Storage


Storage:

Store at -80°C


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84668-2-RR (CEBP Alpha/CEBPA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84668-2-PBS in a different storage buffer




1x10^6 HeLa cells were intracellularly stained with 0.25 ug CEBP Alpha/CEBPA Recombinant antibody (84668-2-RR, Clone:242101C10) and APC-Conjugated Goat Anti-Rabbit IgG(H-L)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and prepabilized with Time. were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. This data was developed using the same antibody clone with 84668-2-PBS in a different storage




1x10^6 HepG2 cells were intracellularly stained with 0.25 ug CEBP Alpha/CEBPA Recombinant antibody (84668-2-RR, Clone:242101C10) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and promobilized with True. were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. This data was developed using the same antibody





Cytometric bead array standard curve of MP01479-1, CEBP Alpha/CEBPA Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84668-1-PBS. Detection antibody: 84668-2-PBS. Standard: Ag29947. Range: 0.625-80 ng/mL



Biolayer interferometry (BLL) kinetic assays of 84668-2-RR against Human CEBP Alpha/CEBPA were performed. The affinity constant is 1.15 nM.