## For Research Use Only

## TFEB Recombinant antibody

Catalog Number:84675-5-RR



**Basic Information** 

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purfication

84675-5-RR Size:

GeneID (NCBI):

otein A purfication

1000 ug/ml

7942

BC032448

CloneNo.: 242093D11

WB 1:500-1:2000

Source:

UNIPROT ID: P19484 Recommended Dilutions:

Rabbit Isotype:

Full Name: transcription factor EB

IgG

Calculated MW:

Immunogen Catalog Number: Ca AG4194 470

476 aa, 53 kDa

Observed MW:

65-70 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Positive Controls:

Species Specificity:

human, mouse

WB: Raji cells, RAW 264.7 cells

## **Background Information**

Transcription factor EB(TFEB) is one of the transcription factors that contain a basic HLH-ZIP domain(for DNA binding ) and HLH and ZIP domains( for loigomerization). TFEB specifically recognizes and binds to E-box sequences, which required dimerization with itself or another MiT/TFE family member. It plays a role in T-cell-dependent antibody responses to activated CD4+ T-cells and thymus-dependent humoral immunity once associated with TFE3. It also activates lysosomal gene by binding the CLEAR-box sequence in the regulatory region. It involves in the signal transduction processes required for normal vascularization of the placenta. This is a rabbit polyclonal antibody raised against the full length of human TFEB. The calculated molecular weight of TFEB is 53 kDa, but modified TFEB is about 65-70 kDa (PMID: 12837690). TFEB mainly present in the cytoplasm (PubMed:23434374). Under aberrant lysosomal storage conditions, it translocates from the cytoplasm to the nucleus (PubMed:23434374). Fisetin increases the levels of TFEB localized in nuclei (PMID: 27112200).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84675-5-RR (TFEB antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 84675-5-RR against Human TFEB were performed. The affinity constant is 7.61 nM.