For Research Use Only

HMGCL Recombinant antibody, PBS Only




Catalog Number:84769-4-PBS


	BC010570	Protein A purfication
Size: 1 mg/ml	GenelD (NCBI): 3155	CloneNo.: 242224F9
Source: Rabbit	UNIPROT ID: P35914	
Isotype: IgG Immunogen Catalog Number:	Full Name: 3-hydroxymethyl-3-methylglutaryl- Coenzyme A lyase	
AG10426	Calculated MW: 325 aa, 34 kDa	
	Observed MW: 32 kDa	
Tested Applications: WB, FC (Intra), Indirect ELISA		
Species Specificity: human, mouse, rat		
HMGCL (3-hydroxy-3-methylglutaryl-CoA lyase) is a key enzyme in ketogenesis (ketone body formation). It is a ketogenic enzyme in the liver that catalyzes the formation of acetoacetate from HMG-CoA within the mitochondria. It also plays a prominent role in the catabolism of the amino acid leucine.		
Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	y at −80°C
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG10426 Tested Applications: WB, FC (Intra), Indirect ELISA Species Specificity: human, mouse, rat HMGCL (3-hydroxy-3-methylgluta ketogenic enzyme in the liver tha It also plays a prominent role in th Storage: Stora at -80°C. The product is shipped with ice pa Storage Buffer:	1 mg/ml 3155 Source: UNIPROT ID: Rabbit P35914 Isotype: Full Name: IgG 3-hydroxymethyl-3-methylglutar Coenzyme A lyase Coenzyme A lyase AG10426 Calculated MW: 325 aa, 34 kDa Observed MW: 32 kDa Observed MW: 32 kDa Species Specificity: human, mouse, rat HMGCL (3-hydroxy-3-methylglutaryl-CoA lyase) is a key enzyme in keth Ketogenic enzyme in the liver that catalyzes the formation of acetoaceth It also plays a prominent role in the catabolism of the amino acid leucint Storage: Storage: Storage Buffer: Storage buffer:

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 84769-4-RR (HMGCL antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84769-4-PBS in a different storage buffer formulation. 1x10^{^6} HeLa cells were intracellularly stained with 0.25 ug HMGCL Recombinant antibody (84769-4-RR, Clone:242224F9) and APC-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84769-4-PBS in a different storage buffer



Biolayer interferometry (BLL) kinetic assays of 84769-4-RR against Human HMGCL were performed. The affinity constant is 1.13 nM.