

For Research Use Only

RNF2 Recombinant antibody, PBS Only

Catalog Number: 84793-5-PBS

Featured Product



Basic Information

Catalog Number:

84793-5-PBS

Size:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG8900

GenBank Accession Number:

BC012583

GeneID (NCBI):

6045

UNIPROT ID:

Q99496

Full Name:

ring finger protein 2

Calculated MW:

336 aa, 38 kDa

Observed MW:

38 kDa

Purification Method:

Protein A purification

CloneNo.:

242181G7

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse

Background Information

RNF2 (also known as ding, Ring1B or Ring2) is a member of the Ring finger protein family, which functions as E3 ubiquitin ligase for monoubiquitination of histone H2A at lysine 119 (H2AK119ub). Many studies have demonstrated that overexpressed RNF2 was involved in the pathological progression of multiple cancers and has an impact on their clinical features. For instance, the upregulated expression level of RNF2 is positively correlated with the occurrence and progression of hepatocellular carcinoma, melanoma, prostate cancer, breast cancer, pancreatic cancer, gastric cancer, and bladder urothelial carcinoma, as well as with the radioresistance of lung cancer and chemoresistance of ovarian cancer (PMID: 33400276). The molecular weight of RNF2 is 38 kDa.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84793-5-RR (RNF2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84793-5-PBS in a different storage buffer formulation.




WB result of RNF2 antibody (84793-5-RR; 1:100000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RNF2 transfected HEK-293 cells. This data was developed using the same antibody clone with 84793-5-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using RNF2 antibody (84793-5-RR, Clone: 242181G7) at dilution of 1:500 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84793-5-PBS in a different storage buffer formulation.



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug RNF2 Recombinant antibody (84793-5-RR, Clone:242181G7) and APC-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. This data was developed using the same antibody clone with 84793-5-PBS in a different storage buffer



Biolayer interferometry (BLI) kinetic assays of 84793-5-RR against Human RNF2 were performed. The affinity constant is 0.251 nM.