For Research Use Only

GPD1 Recombinant antibody

Catalog Number:84800-1-RR



Basic Information

Catalog Number:

GenBank Accession Number: BC032234

Purification Method:

84800-1-RR

Protein A purification

Size:

GeneID (NCBI):

CloneNo.:

1000 µg/ml

2819

242359D3

IF/ICC 1:50-1:500

Source: Rabbit

UNIPROT ID: P21695

Recommended Dilutions: WB 1:500-1:2000

Isotype:

Full Name: glycerol-3-phosphate dehydrogenase

32 kDa

Immunogen Catalog Number:

1 (soluble)

AG4278

Calculated MW: 349 aa, 38 kDa

Observed MW:

Applications

Tested Applications:

Positive Controls:

WB, IF/ICC, ELISA

WB: HepG2 cells, mouse liver tissue

Species Specificity: human, mouse

IF/ICC: HepG2 cells,

Background Information

GPD1 (Glycerol-3-phosphate dehydrogenase 1) is an important enzyme belonging to the NAD-dependent glycerol-3 $phosphate\ dehydrogen as e\ family.\ Its\ C-terminal\ structural\ domain\ contains\ multiple\ helical\ structures\ for\ binding$ the substrate DHAP, and its N-terminal structural domain contains a $\,\beta$ -folded core for binding NADH. GPD1 catalyzes the conversion of dihydroxyacetone phosphate (DHAP) and reduced nicotinamide adenine dinucleotide (NADH) to glycerol-3-phosphate (G3P) and NAD+, and plays a key role in carbohydrate and lipid metabolism. GPD1 also works with mitochondrial glycerol-3-phosphate dehydrogenase to form a glycerophosphate shuttle system that facilitates the transfer of reducing equivalents from the cytoplasm to the mitochondria. Abnormal activity of GPD1 has been associated with a variety of metabolic disorders, such as obesity, hypertriglyceridemia, and GPD1 has been implicated in cancer, potentially acting as a tumor suppressor.


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84800-1-RR (GPD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using GPD1 antibody (84800-1-RR, Clone: 242359D3) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLI) kinetic assays of 84800-1-RR against Human GPD1 were performed. The affinity constant is below 1 pM.