For Research Use Only

BCL3 Recombinant antibody

Catalog Number:84882-5-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

CloneNo.:

Positive Controls:

IHC: mouse spleen tissue,

IF/ICC: HeLa cells, HepG2 cells

242233B11

IHC 1:50-1:500 IF/ICC 1:200-1:800

Basic Information

Catalog Number: GenBank Accession Number: 84882-5-RR BC064993

 Size:
 GeneI D (NCBI):

 1000 μ g/ml
 602

Source: UNIPROT ID:
Rabbit P20749

Isotype: Full Name:

IgG B-cell CLL/lymphoma 3

Immunogen Catalog Number: Calculated MW:
AG13375 446 aa, 47 kDa
Observed MW:

48-60 kDa

Applications

Tested Applications: IHC, IF/ICC, ELISA Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

BCL3 (BCL3 Transcription Coactivator), as a Protein Coding gene, has seven ankyrin repeat domains, a proline-rich N terminal domain and a serine- and proline-rich C-terminal domain(PMID: 31930327). BCL3 is an important contributor to chronic inflammatory disease states, such as osteoarthritis and rheumatoid arthritis. In the cytoplasm, BCL3 inhibits the nuclear translocation of the NF-kappa-B p50 subunit. In the nucleus, BCL3 acts as transcriptional activator that promotes transcription of NF-kappa-B target genes(Uniprot).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 84882-5-RR (BCL3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 84882-5-RR (BCL3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using BCL3 antibody (84882-5-RR, Clone: 242233B11) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 84882-5-RR against Human BCL3 were performed. The affinity constant is 0.581 nM.