For Research Use Only

PTK7 Recombinant antibody

Catalog Number:85075-1-RR





Basic Information	Catalog Number: 85075-1-RR	GenBank Accession Number: BC071557	Purification Method: Protein A purification
	Concentration: 1000 ug/ml	GenelD (NCBI): 5754	CloneNo.: 242720A5
	Source: Rabbit	UNIPROT ID: Q13308	Recommended Dilutions: WB 1:2000-1:10000
	Isotype: IgG	Full Name: PTK7 protein tyrosine kinase 7	
	Immunogen Catalog Number: AG12026	Calculated MW: 1070 aa, 118 kDa	
		Observed MW: 140 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, ELISA Species Specificity: human, mouse, rat	WB : Jurka	at cells, C2C12 cells, PC-12 cells
Background Information	Protein tyrosine kinase 7 (PTK7), also known as colon carcinoma kinase 4 (CCK4), is an atypical receptor tyrosine kinase that consists of an extracellular domain with seven immunoglobulin-like domains, a transmembrane domain, and a catalytically defective tyrosine kinase domain (PMID: 29867084). It is implicated in planar cell polarity and the Wnt canonical and non-canonical pathways (PMID: 24618420). Overexpression of PTK7 has been reported in a variety of tumors, such as cervical cancer, colon cancer, lung adenocarcinoma, acute myelogenous leukemia, and gastric cancer (PMID: 30944666).		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 85075-1-RR (PTK7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.

Various lysates were subjected to SDS PAGE followed by western blot with 85075-1-RR (PTK7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 85075-1-RR against Human PTK7 were performed. The affinity constant is 2.60 nM.