For Research Use Only

KIR2DL1 Recombinant antibody

Catalog Number:85168-1-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

CloneNo.:

242070D11

WB 1:500-1:2000

Basic Information

Catalog Number: 85168-1-RR Concentration:

1000 ug/ml Source: Rabbit Isotype:

cytoplasmic tail, 1 Calculated MW: 39 kDa

45 kDa

Applications

Tested Applications: WB, ELISA

Species Specificity:

human

GenBank Accession Number:

BC069344 GeneID (NCBI): 3802

UNIPROT ID: P43626 Full Name:

killer cell immunoglobulin-like receptor, two domains, long

Observed MW:

Positive Controls:

WB: Recombinant protein,

Background Information

Killer cell immunoglobulin-like receptor 2DL1 (KIR2DL1, also known as CD158A and NKAT1) is an inhibitory receptor of NK cells, with $\,\beta$ -arrestin 2 mediated recruitment of the tyrosine phosphatases SHP-1 and SHP-2 to KIR2DL1 and facilitated downstream inhibitory signaling (PMID: 18604210). The effector function of NK cells expressing KIR2DL1 $or \, KIR2DS1 \, is \, highly \, influenced \, by \, genetic \, variability \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, NK \, receptors \, and \, is \, calibrated \, by \, co-expression \, of \, additional \, of \, calibrated \, and \, calibrated$ and cognate HLA-C2 ligands (PMID: 31024561).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85168-1-RR (KIR2DL1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLL) kinetic assays of 85168-1-RR against Human KIR2DL1 were performed. The affinity constant is 37.9 nM.