

For Research Use Only

DARS2 Recombinant antibody

Catalog Number: 85691-2-RR



Basic Information

Catalog Number:

85691-2-RR

Concentration:

1000 μ g/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4809

GenBank Accession Number:

BC045173

GeneID (NCBI):

55157

UNIPROT ID:

Q6PI48

Full Name:

aspartyl-tRNA synthetase 2,
mitochondrial

Calculated MW:

645 aa, 74 kDa

Observed MW:

66-74 kDa

Purification Method:

Protein A purification

CloneNo.:

243050E5

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : MCF-7 cells, Jurkat cells, LNCaP cells, A549 cells,
HepG2 cells, NCI-H1299 cells

Background Information

DARS2 is also named as AspRS (aspartyl-tRNA synthetase) and belongs to the class-II aminoacyl-tRNA synthetase family. The deduced 645-amino acid protein has a 47-amino acid mitochondrial targeting signal, resulting in a mature protein of 598 amino acids. DARS2 contains conserved residues involved in ATP binding, tRNA binding, and aspartic acid recognition, as well as catalytic site motifs characteristic of amino acid tRNA synthetases. It is a dimeric proteins(PMID:19443655). Rat aspartyl-tRNA synthetase has a N-terminal polypeptide extension of about 40 amino acid residues which can be removed without impairing its catalytic activity(PMID:9030747). This protein has different molecular mass(55 kDa, 50 kDa, 66 kDa) according to the publications(PMID:11306575; 19443655;15299749).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85691-2-RR (DARS2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.