

For Research Use Only

Anti-Human TLR3/CD283 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98016-1-PBS



Basic Information

Catalog Number: 98016-1-PBS	GenBank Accession Number: BC094737	Purification Method: Protein A purification
Size: 1mg, 2 mg/ml	GeneID (NCBI): 7098	CloneNo.: 230445F1
Source: Rabbit	UNIPROT ID: O15455	
Isotype: IgG	Full Name: toll-like receptor 3	
Immunogen Catalog Number: AG34044	Calculated MW: 904 aa, 104 kDa	

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human

Background Information

TLR3 belongs to the Toll-like receptor family which is important in the innate immune response to pathogens. TLRs are highly conserved from *Drosophila* to humans and share structural and functional similarities. TLR3 is a nucleotide-sensing TLR that is activated by double-stranded RNA. Upon recognition of dsRNA, TLR3 transmits signals via the adaptor protein TICAM-1/TRIF, leading to the induction of type I IFN, cytokine/chemokine production, and dendritic cell (DC) maturation (PMID: 18262679).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ human monocyte-derived immature dendritic cells were intracellularly stained with 0.25 ug Anti-Human TLR3/CD283 Rabbit Recombinant Antibody (98016-1-RR, Clone: 230445F1) or 0.25 ug Isotype Control and PE-conjugated Goat Anti-Rabbit IgG. Cells were then stained with APC anti-human CD11b. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98016-1-



1x10⁶ human monocyte-derived immature dendritic cells were intracellularly stained with 0.25 ug Anti-Human TLR3/CD283 Rabbit Recombinant Antibody (98016-1-RR, Clone: 230445F1) (red) or 0.25 ug Isotype Control (blue) and PE-conjugated Goat Anti-Rabbit IgG. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98016-1-PBS in a different storage buffer formulation.



Bi-layer interferometry (BLI) kinetic assays of 98016-1-RR against Human TLR3/CD283 were performed. The affinity constant is 5.56 nM.