

For Research Use Only

# Anti-Human TLR3/CD283 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98016-2-PBS



## Basic Information

<b>Catalog Number:</b> 98016-2-PBS	<b>GenBank Accession Number:</b> BC094737	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1mg, 2 mg/ml	<b>GeneID (NCBI):</b> 7098	<b>CloneNo.:</b> 230445G9
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> O15455	
<b>Isotype:</b> IgG	<b>Full Name:</b> toll-like receptor 3	
<b>Immunogen Catalog Number:</b> AG34044	<b>Calculated MW:</b> 904 aa, 104 kDa	

## Applications

**Tested Applications:**  
FC (Intra)

**Species Specificity:**  
human

## Background Information

TLR3 belongs to the Toll-like receptor family which is important in the innate immune response to pathogens. TLRs are highly conserved from *Drosophila* to humans and share structural and functional similarities. TLR3 is a nucleotide-sensing TLR that is activated by double-stranded RNA. Upon recognition of dsRNA, TLR3 transmits signals via the adaptor protein TICAM-1/TRIF, leading to the induction of type I IFN, cytokine/chemokine production, and dendritic cell (DC) maturation (PMID: 18262679).

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



1x10<sup>6</sup> human monocyte-derived immature dendritic cells were intracellularly stained with 0.25 ug Anti-Human TLR3/CD283 Rabbit Recombinant Antibody (98016-2-RR, Clone: 230445G9) or 0.25 ug Isotype Control and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human CD11c. Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm



1x10<sup>6</sup> human monocyte-derived immature dendritic cells were intracellularly stained with 0.25 ug Anti-Human TLR3/CD283 Rabbit Recombinant Antibody (98016-2-RR, Clone: 230445G9) (red) or 0.25 ug Isotype Control (blue) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human CD11c. Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry