For Research Use Only

## Anti-Human TNF-alpha Rabbit Recombinant Antibody, PBS Only

www.ptglab.com

**Purification Method:** 

Protein A purfication

CloneNo.:

240668B11

Catalog Number: 98064-1-PBS

**Basic Information** 

Catalog Number:

98064-1-PBS

Size:

1mg, 2mg/ml

Source: Rabbit Isotype:

GenBank Accession Number:

NM\_000594.4 GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000232810

**UNIPROT ID:** P01375 Full Name:

tumor necrosis factor (TNF

superfamily, member 2)

Calculated MW: 26 kDa

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human

**Background Information** 

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, ins resistance, and cancer.

Storage


Storage:

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:


PBS Only

## **Selected Validation Data**






1x10^6 untreated or PMA, lonomycin and Brefeldin A treated human PBMCs were intracellularly stained with 0.25 ug Anti-Human TNF-alpha Rabbit Recombinant Antibody (98064-1-RR, Clone:240668B11) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)Cells were then stained with CoraLite® Plus 647 Anti-Human CD4. Cells were fixed with 4% PFA and permeabilized with Intracellular Staining Permeabilization Wash Buffer. This data was developed using



1x10^6 untreated(black) or PMA, Ionomycin and Brefeldin A treated(red) human PBMCs were intracellularly stained with 0.25 ug Anti-Human TNF-alpha Rabbit Recombinant Antibody (98064-1-RR, Clone:240668B11) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), or 0.25 ug Isotype Control(blue). Cells were fixed with 4% PFA and permeabilized with Intracellular Staining Permeabilization Wash Buffer. This data was developed using the same antibody clone



Biolayer interferometry (BLI) kinetic assays of 98064-1-RR against Human TNF-alpha were performed. The affinity constant is below 1 pM.