For Research Use Only

Anti-Human TNF-alpha Rabbit Recombinant Antibody

Catalog Number: 98064-1-RR



Basic Information

Catalog Number:

98064-1-RR

Size:

100ug, 1000 $\,\mu$ g/ml

Source: Rabbit Isotype:

i i

UNIPROT ID:
P01375
Full Name:
tumor necrosis factor (TNF

superfamily, member 2)
Calculated MW:

GenBank Accession Number:

NM_000594.4

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000232810

26 kDa

Purification Method: Protein A purfication

CloneNo.: 240668B11

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Background Information

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, ins resistance, and cancer.

Storage


Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH 7.3.

Selected Validation Data






1x10^6 untreated or PMA, Ionomycin and Brefeldin A treated human PBMCs were intracellularly stained with 0.25 ug Anti-Human TNF-alpha Rabbit Recombinant Antibody (98064-1-RR, Clone:240668B11) and PE-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L)Cells were then stained with Coralite® Plus 647 Anti-Human CD4. Cells were fixed with 4% PFA and permeabilized with Intracellular Staining Permeabilization Wash Buffer.



1x10^6 untreated(black) or PMA, Ionomycin and Brefeldin A treated(red) human PBMCs were intracellularly stained with 0.25 ug Anti-Human TNF-alpha Rabbit Recombinant Antibody (98064-1-RR, Clone:240668B11) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), or 0.25 ug Isotype Control(blue). Cells were fixed with 4% PFA and permeabilized with Intracellular Staining Permeabilization Wash Buffer.



Biolayer interferometry (BLI) kinetic assays of 98064-1-RR against Human TNF-alpha were performed. The affinity constant is below 1 pM.