

For Research Use Only

# Anti-Human SIRP Alpha/CD172a Rabbit Recombinant Antibody

Catalog Number: 98101-1-RR



## Basic Information

Catalog Number:

98101-1-RR

Size:

100ug, 1000 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM\_080792

GeneID (NCBI):

140885

UNIPROT ID:

P78324

Full Name:

signal-regulatory protein alpha

Calculated MW:

55 kDa

Purification Method:

Protein A purification

CloneNo.:

240789E1

## Applications

Tested Applications:

FC

Species Specificity:

human

## Background Information

SIRP Alpha, also known as CD172a, SHPS-1, and BIT, belongs to the SIRP family. SIRP Alpha is a transmembrane glycoprotein expressed explicitly on myeloid cells and provides a "do not eat me" signal after engaging with integrin-associated protein CD47 on tumor cells (PMID: 36419386). SIRP Alpha shows heterogeneity in molecular weight (~65-120 kDa) in various tissues due to differential glycosylation (PMID: 18051954).

## Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH 7.3.

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



$1 \times 10^6$  human peripheral blood leukocytes were surface stained with 0.25  $\mu$ g Anti-Human SIRP Alpha/CD172a Rabbit Recombinant Antibody (98101-1-RR, Clone: 240789E1) or Isotype Control, and PE-conjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



$1 \times 10^6$  human peripheral blood leukocytes were surface stained with 0.25  $\mu$ g Anti-Human SIRPA Rabbit Recombinant Antibody (98101-1-RR, Clone: 240789E1) (red) or Isotype Control (blue), and PE-conjugated Goat Anti-Rabbit IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



Bi-layer interferometry (BLI) kinetic assays of 98101-1-RR against Human SIRP Alpha/CD172a were performed. The affinity constant is 3.39 nM.