For Research Use Only

## Anti-Human IL3RA/CD123 Rabbit Recombinant Antibody, PBS Only

Catalog Number:98107-1-PBS




Basic Information	Catalog Number: 98107-1-PBS	GenBank Accession Number: BC035407	Purification Method: Protein A purfication	
	Size: 1mg, 2mg/ml	GenelD (NCBI): 3563	CloneNo.: 241268E3	
	Source: Rabbit	ENSEMBL Gene ID: ENSG0000185291		
	Isotype: IgG	UNIPROT ID: P26951		
		Full Name: interleukin 3 receptor, alpha (low affinity)	,	
		Calculated MW: 378 aa, 43 kDa		
Applications	Tested Applications: FC			
	Species Specificity: human			
Background Information	glycoprotein containing an e transmembrane domain and progenitor cells, plasmacyto malignancies (PMID: 105274 (CD131) to form the high-aff JAK/STAT, Ras-MAPK, and ph	Interleukin-3 receptor subunit alpha (IL3RA), also known as CD123, belongs to the cytokine receptor family. It is a glycoprotein containing an extracellular domain, involving a predicted Ig-like domain, two FnIII domains, a transmembrane domain and an intracellular domain (PMID: 24513123). IL3RA is expressed on hematopoietic progenitor cells, plasmacytoid dendritic cells, basophils, monocytes, some B cells, and in a variety of hematologic malignancies (PMID: 10527461; 34855461). IL-3 initially binds to IL3RA and subsequently recruits the beta chain (CD131) to form the high-affinity receptor, resulting in the activation of a series of signaling pathways including the JAK/STAT, Ras-MAPK, and phosphatidylinositol 3-kinase pathways (PMID: 29296749). IL-3R signaling regulates the proliferation, survival, and differentiation of hematopoietic cells.		
Storage	Storage: Store at -80°C. The product is shipped with i Storage Buffer: PBS Only	ice packs. Upon receipt, store it immediatel	y at −80°C	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>^6</sup> human PBMCs were surface stained with 0.25 ug Anti-Human IL3RA/CD123 Rabbit Recombinant Antibody (98107-1-RR, Clone: 241268E3) or 0.25 ug Isotype Control and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG (H+L). Cells were co-stained with Coralite® Plus 647 Anti-Human CD45R0 (UCHL1) (CL647-65150, Clone: UCHL1). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. Lymphocytes were gated. This data was developed using