

For Research Use Only

Anti-Human CD84 Rabbit Recombinant Antibody

Catalog Number: 98108-1-RR



Basic Information

Catalog Number: 98108-1-RR	GenBank Accession Number: BC020063	Purification Method: Protein A purification
Size: 100ug, 1000 ug/ml	GeneID (NCBI): 8832	CloneNo.: 241345A3
Source: Rabbit	UNIPROT ID: Q9UIB8	
Isotype: IgG	Full Name: CD84 molecule	
	Calculated MW: 345 aa, 39 kDa	

Applications

Tested Applications:
FC

Species Specificity:
human

Background Information

CD84, also known as SLAMF5, is a member of the CD2 subfamily of the immunoglobulin receptor superfamily (PMID: 9310491). CD84 is a single-chain type-I glycoprotein composed of two extracellular Ig-like domains, a hydrophobic transmembrane region, and a cytoplasmic domain. It is broadly expressed on almost all leukocyte subsets. CD84 functions as a homophilic adhesion molecule, whose signaling can activate or inhibit leukocyte function depending on the cell type and its stage of activation or differentiation (PMID: 30522694).

Storage

Storage:
Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide, pH 7.3.

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD84 Rabbit Recombinant Antibody (98108-1-RR, Clone: 241345A3) or Isotype Control, and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were then stained with CoraLite® Plus 647 Anti-Human CD4 (OKT4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD84 Rabbit Recombinant Antibody (98108-1-RR, Clone: 241345A3) or Isotype Control, and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



Biolayer interferometry (BLI) kinetic assays of 98108-1-RR against Human CD84 were performed. The affinity constant is 1.14 nM.