For Research Use Only

## Anti-Mouse F4/80 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98236-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

98236-1-PBS

NM\_010130.4 GeneID (NCBI): Protein A purfication

Size:

CloneNo.:

1mg, 2 mg/ml Source:

**UNIPROT ID:** 

241959G4

Rabbit Isotype: Q61549 Full Name:

EGF-like module containing, mucin-

like, hormone receptor-like sequence

Calculated MW:

102 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

mouse

**Background Information** 

Mouse F4/80, also named as EMR1 and Gpf480, is a 160-kDa cell surface glycoprotein which is a member of the EGF TM7 family. The F4/80 molecule is solely expressed on the surface of macrophages and serves as a marker for mature macrophage tissues, including Kupffer cells in liver, splenic red pulp macrophages, brain microglia, gut lamina propria and Langerhans cells in the skin. The function of F4/80 is unclear, but it is speculated to be involved in macrophage adhesion events, cell migration or as a G protein-coupled signaling component of macrophages.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only

## **Selected Validation Data**



1x10^6 Balb/c mouse peritoneal macrophages were surface stained with 0.25 ug Anti-Mouse F4/80 Rabbit Recombinant Antibody (98236-1-RR, Clone: 241959G4) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with CoraLite® Plus 647 Anti-Mouse CD11b. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data



1x10^6 Balb/c mouse peritoneal macrophages were surface stained with 0.25 ug Anti-Mouse F4/80 Rabbit Recombinant Antibody (98236-1-RR, Clone: 241959G4) (red) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98236-1-PBS in