FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

Recombinant human PTPN14 protein



Basic Information	Catalog Number:	Peptide Sequence:		
	Ag12571	MPFGLKLRRTRRYNVLSKNCFVTRIRLLDSNVIECTLS		
	Size:	ESTGQECLEAVAQRLELRETHYFGLWFLSKSQQARW VELEKPLKKHLDKFANEPLLFFGVMFYVPNVSWLQQI		
	50 µg	ATRYQYYLQVKKDVLEGRLRCTLDQVIRLAGLAVQAI		
	Form: FGDYNQFDSQDFLREYVLFPMDLALEE Available lyophilized AQEHKAHSGILPAEAELMYINEVERLDG Species: DNHGNCVHLGIFFMGIFVRNRIGRQAV human ITHNKSTILVELINKEETALFHTDDIENA	FGDYNQFDSQDFLREYVLFPMDLALEEAVLEELTQKV AQEHKAHSGILPAEAELMYINEVERLDGFGQEIFPVK		
		DNHGNCVHLGIFFMGIFVRNRIGRQAVIYRWNDMGI ITHNKSTILVELINKEETALFHTDDIENAKYISRLFATR KFYKQNKICTEQSNSPPPIRRQPTWSRSSLPRQQPYI PPVHVQCGEHYSET		
			Endotoxin Level: Please contact the lab for more information	
			Reconstitution	Reconstitution:
	and Storage	Reconstitute at 0.25 µg/ µ l in 200 µ l sterile water for short- term storage. After reconstitution with sterile water, if glycerol has no effect on subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).	
equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the				
ion concentration of the final solution will vary according to				
the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.				
Purity	85%, by SDS-PAGE with Coomassie Brilliant Blue staining.			
Formulation	The purified protein was Lyophilized from sterile PBS (58mM Na2ł trehalose and 5 % mannitol are added as protectant before lyophi	• •		
Stability and Storage	Store for up to 12 months at -20°C to -80°C as lyophilized powder.			
Storage of Reconstituted Protein	Short Term Storage: Store at 2-8°C for (1-2 weeks). Long Term Storage: Aliquot and store at -20°C to -80°C for up to 3 months, buffer			
	containing 50% glycerol is recommended for reconstitution. Avoid repeat freeze-thaw cycles.			
Selected Validation Data				
	76 kDa→ ← 64 kDa			
	43 kDa→			
	28 kDa→			

20 kI