## FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

## Recombinant human USF1 protein



Basic Information	Catalog Number:	Peptide Sequence:
	Ag17789	MKGQQKTAETEEGTVQIQEGAVATGEDPTSVAIASI
	Size:	QSAATFPDPNVKYVFRTENGGQVMYRVIQVSEGQL DGQTEGTGAISGYPATQSMTQAVIQGAFTSDDAVD
	50 µ g	TEGTAAETHYTYFPSTAVGDGAGGTTSGSTAAVVTT
	Form: Available lyophilized	QGSEALLGQATPPGTGQFFVMMSPQEVLQGGSQRS IAPRTHPYSPKSEAPRTTRDEKRRAQHNEVERRRRDKI NNWIVQLSKIIPDCSMESTKSGQSKGGILSKACDYIQE LRQSNHRLSEELQGLDQLQLDNDVLRQQVEDLKNKNL
	Species: human	
	Expression Source: <i>e coli-</i> derived, PGEX-4T, with N-terminal GST.	LLRAQLRHHGLEVVIKNDSN (1-310 aa encoded by BC035505)
	Biological Activity: Not tested	
	Endotoxin Level: Please contact the lab for more information	
Reconstitution and Storage	Reconstitution: Reconstitute at 0.25 µg/ µ l in 200 µ l sterile water for short- term storage. After reconstitution with sterile water, if glycerol has no effect	Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).
	on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please	· · · ·
	adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full	
Purity	recovery of the protein into solution. 85%, by SDS-PAGE with Coomassie Brilliant Blue staining.	
Formulation	The purified protein was Lyophilized from sterile PBS (58mM Na2 trehalose and 5 % mannitol are added as protectant before lyoph	•
Stability and Storage	Store for up to 12 months at -20°C to -80°C as lyophilized powder.	
Storage of Reconstituted Protein	Short Term Storage: Store at 2-8°C for (1-2 weeks). Long Term Storage: Aliquot and store at -20°C to -80°C for up to 3 months, buffer containing 50% glycerol is recommended for reconstitution. Avoid repeat freeze-thaw cycles.	
Selected Validation Data	74 kDa→ 66 kDa→ ⊷-64 kDa	
	43 kDa→ 28 kDa→	

20 kDa