For Research Use Only

CoraLite® Plus 405 Anti-Human CD86 (BU63) Mouse IgG2a Recombinant Antibody



Catalog Number: CL405-65592

Basic Information

Catalog Number: CL405-65592 Concentration:

100tests, 5 ul/test Source: Mouse

Isotype: IgG2a GenBank Accession Number: BC040261

GenelD (NCBI):

ENSEMBL Gene ID: ENSG00000114013 Full Name: CD86 molecule Calculated MW:

329 aa, 38 kDa

Purification Method: Protein A purification

CloneNo.: BU63

Excitation/Emission maxima wavelengths:
399 nm / 422 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of CD86 with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of CD86 with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 0.09% sodium azide.

Selected Validation Data





1x10^6 human PBMCs were surface stained with APC Anti-Human CD14 Rabbit Recombinant Antibody (APC-98040, Clone: 230332D7), and 5 ul CoraLite® Plus 405 Anti-Human CD86 (BU63) Mouse IgG2a RecAb (CL405-65592, Clone: BU63) or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



CL405-65592 CD86(BU63),PB450-A

1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 405 Anti-Human CD86 (BU63) Mouse IgG2a RecAb (CL405-65592, Clone:BU63) (red) or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.