For Research Use Only

CoraLite® Plus 488-conjugated IRF3 Polyclonal antibody



Purification Method:

IF 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-11312

Featured Product

Basic Information

Applications

Catalog Number: CL488-11312

1000 µg/ml Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG1858

interferon regulatory factor 3 Calculated MW:

Observed MW: 50-60 kDa

Tested Applications: FC (Intra), IF/ICC

Species Specificity:

human

GenBank Accession Number:

BC009395 GeneID (NCBI):

3661 **UNIPROT ID:**

Q14653

47 kDa

Full Name:

Positive Controls: IF: HepG2 cells,

Background Information

The virul-induced expression of interferon(IFN) genes in infected cells implicate in the interplay of several constitutively expressed and virus-activated transcription factors. A family of IFN regulatory factors(IRFs) have been shown to has a role in the transcription of IFN genes as well as IFN-stimulated genes. IRF3 is a novel key transcriptional regulator of type I IFN-dependent immune responses and involves in the innate immune response against DNA and RNA viruses, by binding to the promoters of IFN. It located in the cytoplasm of uninfected cells in an inactive form, and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, could be phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization, nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes. IRF3 exists some isoforms with MV 47-49 kDa, 33 kDa and 12-16 kDa.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 IRF3 antibody (CL488-11312) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human IRF3 (CL488-11312) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).