For Research Use Only

CoraLite® Plus 488-conjugated IRF3 Polyclonal antibody

Catalog Number: CL488-11312

Featured Product

1 Publications

BC009395

3661

GeneID (NCBI):

GenBank Accession Number:



Basic Information

Catalog Number: CL488-11312 Concentration: 1000 ug/ml

Source: UNIPROT ID:
Rabbit Q14653

Isotype: Full Name:
IgG interferon regulatory factor 3

Immunogen Catalog Number: Calculated MW:

AG1858 47 kDa

Observed MW: 50-60 kDa **Purification Method:**

Antigen affinity purification

Recommended Dilutions: IF/ICC: 1:50-1:500

FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF/ICC, FC (Intra)
Cited Applications:

IF

Species Specificity:

human
Cited Species:
human

Positive Controls:

IF/ICC: HepG2 cells,

FC (Intra): human peripheral blood lymphocytes,

Background Information

The virul-induced expression of interferon(IFN) genes in infected cells implicate in the interplay of several constitutively expressed and virus-activated transcription factors. A family of IFN regulatory factors(IRFs) have been shown to has a role in the transcription of IFN genes as well as IFN-stimulated genes. IRF3 is a novel key transcriptional regulator of type I IFN-dependent immune responses and involves in the innate immune response against DNA and RNA viruses, by binding to the promoters of IFN. It located in the cytoplasm of uninfected cells in an inactive form, and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, could be phosphorylated by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization, nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes. IRF3 exists some isoforms with MV 47-49 kDa, 33 kDa and 12-16 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Lin Zhu	38285487	Elife	IF

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 IRF3 antibody (CL488-11312) at dilution of 1:200.



1x10^6 human peripheral blood lymphocytes were intracellularly stained with 0.8 ug Coralite® Plus 488-conjugated IRF3 Polyclonal antibody (CL488-11312)(red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit 1gG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).