For Research Use Only

CoraLite® Plus 488-conjugated ASGR1 Polyclonal antibody



Catalog Number: CL488-11739 1 Publications

Basic Information

Catalog Number: CL488-11739

BC032130

GeneID (NCBI):

GenBank Accession Number:

1000 µ g/ml

UNIPROT ID: P07306

Rabbit P07306 Isotype: Full Name:

gG asialoglycoprotein receptor 1

Immunogen Catalog Number: Calculated MW: AG2310 291 aa, 33 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: IF-P 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

IF-P, FC

Source:

Cited Applications:

IF

Species Specificity: human, mouse, rat Cited Species: mouse Positive Controls:

IF-P: rat liver tissue, human liver cancer tissue

Background Information

Asialoglycoprotein receptor (ASGPR), also known as the hepatic galactose/N-acetylglucosamine (GlcNAc) receptor or Ashwell receptor, is a C-type lectin expressed exclusively in hepatic parenchymal cells. ASGPR consists of two subunits, a major subunit (ASGR1, HL-1) and a minor subunit (ASGR2, HL-2), and specifically recognizes terminal β -linked galactose or GlcNAc on circulating glycoproteins or cells. This receptor plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins that contain terminal galactose or GlcNAc residues. ASGPR may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery. This antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

Notable Publications

Author	Pubmed ID	Journal	Application
Tian Tian	37829506	MedComm (2020)	IF

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite® Plus 488 ASGR1 antibody (CL488-11739) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed rat liver tissue using CoraLite® Plus 488 ASGR1 antibody (CL488-11739) at dilution of 1:200.



1X10^6 HepG2 cells were surface stained with 0.4 ug CoraLite® Plus 488 Anti-Human ASGR1 (CL488-11739) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit 1gG control Rabbit PolyAb (CL488-30000, Clone:) (blue). Cells were not fixed.