For Research Use Only

CoraLite® Plus 488-conjugated TOM20 Polyclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: CL488-11802

Featured Product

4 Publications

Basic Information

Catalog Number: CL488-11802

Size: 1000 µg/ml Source: Rabbit

gG

Immunogen Catalog Number:

AG2378

Isotype:

GenBank Accession Number:

BC000882 GeneID (NCBI): 9804 UNIPROT ID: Q15388 Full Name:

translocase of outer mitochondrial membrane 20 homolog (yeast)

Calculated MW: 145 aa, 16 kDa Observed MW: 16 kDa Purification Method:

Antigen affinity purification
Recommended Dilutions:
IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

493 nm / 522 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Cited Applications:

IF

Species Specificity: human, mouse, rat Cited Species: human, mouse Positive Controls:

IF/ICC: HUVEC cells,

Background Information

TOM20, also named as KIAA0016, belongs to the Tom20 family. It is a central component of the receptor complex responsible for the recognition and translocation of cytosolically synthesized mitochondrial preproteins. Together with TOM22, TOM20 functions as the transit peptide receptor at the surface of the mitochondrion outer membrane and facilitates the movement of preproteins into the TOM40 translocation pore. TOM20 is characterized as major docking receptors to mediate the recognition by different mechanisms.

Notable Publications

Author	Pubmed ID	Journal	Application
Werner J Geldenhuys	34496224	Free Radic Biol Med	IF
Neeraja Purandare	36339251	iScience	IF
Ziqi Guo	39182128	Biol Direct	IF

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using CoraLite® Plus 488 TOM20 antibody (CL488-11802) at dilution of 1:200, CL594-Phalloidin (red).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human TOM20 (CL488-11802) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).