

For Research Use Only

CoraLite® Plus 488-conjugated PGC1a Monoclonal antibody



Catalog Number: CL488-66369

Featured Product

Basic Information

Catalog Number:

CL488-66369

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

NM_013261

GeneID (NCBI):

10891

UNIPROT ID:

Q9UBK2

Full Name:

peroxisome proliferator-activated
receptor gamma, coactivator 1 alpha

Calculated MW:

91 kDa

Purification Method:

Protein G purification

CloneNo.:

1C1B2

Recommended Dilutions:

IF 1:200-1:800

Excitation/Emission maxima
wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

FC (Intra), IF/ICC

Species Specificity:

human, mouse, rat

Positive Controls:

IF : HeLa cells,

Background Information

PPARGC1A, also named as Peroxisome proliferator-activated receptor gamma coactivator 1-alpha, is a 798 amino acid protein, which contains 1 RRM (RNA recognition motif) domain and localizes in the nucleus. PPARGC1A is transcriptional coactivator for steroid receptors and nuclear receptors. PPARGC1A can regulate key mitochondrial genes that contribute to the program of adaptive thermogenesis and plays an essential role in metabolic reprogramming in response to dietary availability through coordination of the expression of a wide array of genes involved in glucose and fatty acid metabolism. PPARGC1A exists as isoforms in the liver and the molecular weight of it is 77 kDa. Except for a deletion of 127 amino acids at the N terminus, the protein, termed L-PGC-1 α , is identical to PGC-1.

Storage

Storage:

Store at -20°C. Avoid exposure to light.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:



T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human PGC1a (CL488-66369, Clone:1C1B2) (red), or 0.2 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite® Plus 488 PGC1a antibody (CL488-66369, Clone: 1C1B2) at dilution of 1:400.