

For Research Use Only

CoraLite® Plus 488-conjugated YAP1 Monoclonal antibody



Catalog Number: CL488-66900

2 Publications

Basic Information

Catalog Number:

CL488-66900

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG28194

GenBank Accession Number:

BC038235

GeneID (NCBI):

10413

UNIPROT ID:

P46937

Full Name:

Yes-associated protein 1, 65kDa

Calculated MW:

504 aa, 54 kDa

Observed MW:

70 kDa

Purification Method:

Protein G purification

CloneNo.:

3A7A9

Recommended Dilutions:

IF 1:10-1:200

**Excitation/Emission maxima
wavelengths:**

493 nm / 522 nm

Applications

Tested Applications:

IF/ICC

Cited Applications:

IF, WB

Species Specificity:

Human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

IF : HepG2 cells,

Background Information

Yes-associated protein 1 (YAP1) is a transcriptional regulator which can act both as a coactivator and a corepressor and is the critical downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Plays a key role to control cell proliferation in response to cell contact. Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. The presence of TEAD transcription factors are required for it to stimulate gene expression, cell growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction. Isoform 2 and isoform 3 can activate the C-terminal fragment (CTF) of ERBB4 (isoform 3). Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level). It is activated by phosphorylation and degraded by ubiquitination (20048001). The calculated molecular weight of YAP1 is 54 kDa, but phosphorylated YAP1 is about 65-70 kDa. (PMID: 26695440)

Notable Publications

Author	Pubmed ID	Journal	Application
Yue Wan	36598105	Glia	WB
Zengshu Huang	36552052	Biomedicines	IF

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using the CoraLite® Plus 488-conjugated version of this antibody, CL488-66900 (YAP1 antibody), at dilution of 1:100.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite® Plus 488 YAP1 antibody (CL488-66900, Clone: 3A7A9) at dilution of 1:200.