For Research Use Only

## CoraLite® Plus 488-conjugated Phospho-mTOR (Ser2448) Monoclonal antibody



Catalog Number: CL 488-67778

**Basic Information** 

Catalog Number: GenBank Accession Number: Purification Method: CL488-67778 BC117166 Protein A purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 1000 μ g/ml
 2475
 2A12G3

Source: Full Name: Recommended Dilutions:

Mouse FK506 binding protein 12-rapamycin IF 1:50-1:500

 Isotype:
 associated protein 1
 Excitation/Emission maxima

 IgG2b
 Calculated MW:
 wavelengths:

Calculated MW: wavelengths: 289 kDa 493 nm / 522 nm

**Applications** 

Tested Applications: Positive Controls: FC (Intra), IF/ICC IF : HepG2 cells, Species Specificity:

## **Background Information**

MTOR, also named as FRAP1, FRAP, FRAP2 and RAPT1, belongs to the PI3/PI4-kinase family. MTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. MTOR is kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTOR is phosphorylated at Ser2448 via the PI3 kinase/Akt signaling pathway and autophosphorylated at Ser2481. mTOR plays a key role in cell growth and homeostasis and may be abnormally regulated in tumors.

Storage

Storage:

Store at -20°C. Avoid exposure to light.


Storage Buffer:

Human, Mouse


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 PhosphomTOR (Ser2448) antibody (CL488-67778, Clone: 2A12G3) at dilution of 1:100. DAPI (blue)



1X10^6 HeLa cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.13 ug Coralite® Plus 488 Anti-Human Phospho-mTOR (Ser2448) (CL488-67778, Clone:2A12G3), or 0.13 ug Control Antibody (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.