For Research Use Only

CoraLite® Plus 488-conjugated EIF2S1 Monoclonal antibody


Catalog Number:CL488-68479 Featured Product

oroteintec Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: CL488-68479	GenBank Accession Number: BC002513	Purification Method: Protein G purification	
	Size: 1000 μg/ml	GenelD (NCBI): 1965	CloneNo.: 3C8E2	
	Source: Mouse	UNIPROT ID: P05198	Recommended Dilutions: IF 1:50-1:500	
	Isotype: IgG1 Immunogen Catalog Number: AG28309	Full Name: eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa Calculated MW: 36 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm	
				Observed MW: 36 kDa
		Applications		Tested Applications: IF/ICC
Species Specificity: Human, Mouse, Rat				
		EIF2S1 is one subunit of the translation initiation factor EIF2, which catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B. EIF2A (Gene ID: 83939) and EIF2S1 (Gene ID: 1965) share the EIF2A symbol/alias in common. EIF2S1 is the alpha subunit of the eIF2 translation initiation complex. Although both of these proteins function in binding initiator tRNA to the 40S ribosomal subunit, the EIF2A protein does so in a codon-dependent manner, whereas eIF2 complex requires GTP.		
Background Information	synthesis initiation, promoting the 40S ribosomal subunit, followed be ribosomal subunit to form the 80S release of an eIF-2-GDP binary co GDP bound to eIF-2 must exchang EIF2S1 (Gene ID: 1965) share the be initiation complex. Although both	e binding of the initiator tRNA to 40S rib by mRNA binding to form a 43S preinitia 5 initiation complex is preceded by hydr mplex. In order for eIF-2 to recycle and ge with GTP by way of a reaction catalyz EIF2A symbol/alias in common. EIF2S1 n of these proteins function in binding in	osomal subunits. This complex binds to a tion complex. Junction of the 60S olysis of the GTP bound to eIF-2 and catalyze another round of initiation, the ed by eIF-2B. EIF2A (Gene ID: 83939) and is the alpha subunit of the eIF2 translation itiator tRNA to the 40S ribosomal subunit,	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 EIF2S1 antibody (CL488-68479, Clone: 3C8E2) at dilution of 1:200, CL594-Phalloidin (red).