## For Research Use Only

## CoraLite® Plus 488-conjugated ADRP Recombinant antibody

Catalog Number:CL488-80362-2


Basic Information	Catalog Number: CL488-80362-2	GenBank Accession Number: BC005127	Purification Method: Protein A purification
	Size: 1000 µg/ml	GenelD (NCBI): 123	CloneNo.: 230274F4
	Source: Rabbit	UNIPROT ID: Q99541	Recommended Dilutions: IF/ICC 1:50-1:500
	Isotype: IgG Immunogen Catalog Number: AG7539	Full Name: adipose differentiation-related protein	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
		Calculated MW: 48 kDa	
		Observed MW: 48 kDa	
Applications	Tested Applications: IF/ICC, FC (Intra)	Positive Controls:	
	IF/ICC : oleic acid treated HeLa cells, Species Specificity: human		
Background Information	ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed oleic acid treated HeLa cells using CoraLite® Plus 488 ADRP antibody (CL488-80362-2, Clone: 230274F4) at dilution of 1:200, CL594-phalloidin (red).



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite® Plus 488 ADRP Recombinant Antibody (CL488-80362-2, Clone:230274F4) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.8 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).